

DESCRIPTION

LabDiet® JL 6% Oval 5K0Q is an autoclavable, complete life-cycle diet formulated using managed formulation, Constant Nutrition®. LabDiet® 5K0Q is the primary diet used at The Jackson Laboratory. This diet does not contain silicon dioxide, an additive which helps to reduce diet clumping post-autoclaving.

Features and Benefits

- Excellent diet to meet the nutrient requirements for maintenance, breeding and lactation.
- 5K0Q is a primary diet used at The Jackson Laboratory. Specific information on strains fed can be obtained from The Jackson Laboratory.
- Standard oval shaped feed pellet.
- Does not contain silicon dioxide, an additive used to reduce diet clumping post-autoclaving.
- Fortified with extra nutrients to compensate for losses during processing and autoclaving.
- If a cylinder shaped pellet is desired, please refer to 5K52.

Product Forms Available

- Oval shaped - 3/8" x 5/8" x 1" length

Other Versions Available

- 5K52: Cylinder shaped pellet - 3/8" diameter by 3/4" length
- 5LG4/5LL4 JL Rat and Mouse Irr 6F

GUARANTEED ANALYSIS

Crude protein not less than	18.0%
Crude fat not less than	6.0%
Crude fiber not more than	5.0%
Ash not more than	8.0%

INGREDIENTS

Ground wheat, ground corn, wheat middlings, ground oats, fish meal, dehulled soybean meal, soybean oil, corn gluten meal, dehydrated alfalfa meal, dicalcium phosphate, monocalcium phosphate, brewers dried yeast, calcium carbonate, menadione dimethylpyrimidinol bisulfite, salt, DL-methionine, choline chloride, magnesium oxide, thiamin mononitrate, pyridoxine hydrochloride, cholecalciferol, vitamin A acetate, calcium pantothenate, ferrous sulfate, biotin, manganous oxide, dl-alpha tocopheryl acetate, folic acid, vitamin B₁₂ supplement, riboflavin, nicotinic acid, zinc oxide, ferrous carbonate, copper sulfate, zinc sulfate, cobalt carbonate, calcium iodate.

FEEDING DIRECTIONS

Feed ad libitum to rodents. Provide plenty of fresh clean water at all times.

AUTOCCLAVING SUGGESTIONS

During the autoclaving process, the pellets can be placed on trays, in small bags or in larger bags, as long as the pellets are stacked no more than 3 inches high. When steam autoclaved, the pellets swell and exert force on adjacent pellets. If confined by a bag or container, the pressure causes sticking as greater polymerization of fibrous materials occurs under such conditions. **Assay before and after autoclaving:** Conditions of sterilization must be determined for each autoclaving unit. It is best to assay the diet before and after sterilization to determine nutrient losses. Microbiological studies should be done also to insure the degree of sterilization desired.

For Product Availability, visit www.labdiet.com.

CHEMICAL COMPOSITION

Nutrients**

Protein, %	19.3
Arginine, %	1.03
Cystine, %	0.25
Glycine, %	0.94
Histidine, %	0.44
Isoleucine, %	0.87
Leucine, %	1.52
Lysine, %	0.97
Methionine, %	0.73
Phenylalanine, %	0.85
Tyrosine, %	0.56
Threonine, %	0.68
Tryptophan, %	0.23
Valine, %	0.90
Serine, %	0.98
Aspartic Acid, %	1.87
Glutamic Acid, %	4.52
Alanine, %	1.13
Proline, %	1.53
Taurine, %	0.03

Fat (ether extract), % .62

Fat (acid hydrolysis), % .72

Cholesterol, ppm	240
Linoleic Acid, %	2.88
Linolenic Acid, %	0.37
Arachidonic Acid, %	0.01
Omega-3 Fatty Acids, %	0.46
Total Saturated Fatty Acids, %	1.24
Total Monosaturated	
Fatty Acids, %	1.37

Fiber (Crude), % .43

Neutral Detergent Fiber³, % .15.1

Acid Detergent Fiber⁴, % .5.2

Nitrogen-Free Extract

(by difference), % .53.6

Starch, % .38.9

Glucose, % .0.12

Fructose, % .0.15

Sucrose, % .0.62

Lactose, % .0.00

Total Digestible Nutrients, % .76.3

Gross Energy, kcal/gm .4.17

Physiological Fuel Value⁵,

kcal/gm .3.47

Metabolizable Energy,

kcal/gm .3.17

Minerals

Ash, % .6.5

Calcium, % .1.17

Phosphorus, % .0.93

Phosphorus (non-phytate), % .0.68

Potassium, % .0.66

Magnesium, % .0.22

Sulfur, % .0.33

Sodium, % .0.26

Chlorine, % .0.45

Fluorine, ppm .37

Iron, ppm .380

Zinc, ppm .85

Manganese, ppm .160

Copper, ppm .11

Cobalt, ppm .0.80

Iodine, ppm .2.1

Chromium, ppm .2.0

Selenium, ppm .0.30

Vitamins

Carotene, ppm .1.5

Vitamin K (as menadione), ppm .20

Thiamin Hydrochloride, ppm .79

Riboflavin, ppm .9.0

Niacin, ppm .90

Pantothenic Acid, ppm .37

Choline Chloride, ppm .2000

Folic Acid, ppm .1.9

Pyridoxine, ppm .10

Biotin, ppm .0.30

B₁₂, mcg/kg .50

Vitamin A, IU/gm .20

Vitamin D₃ (added), IU/gm .4.3

Vitamin E, IU/kg .45

Ascorbic Acid, mg/gm .—

Calories provided by:

Protein, % .22.238

Fat (ether extract), % .16.028

Carbohydrates, % .61.734

*Product Code

1. Formulation based on calculated values from the latest ingredient analysis information. Since nutrient composition of natural ingredients varies and some nutrient loss will occur due to manufacturing processes, analysis will differ accordingly.
2. Nutrients expressed as percent of ration except where otherwise indicated. Moisture content is assumed to be 10.0% for the purpose of calculations.
3. NDF = approximately cellulose, hemi-cellulose and lignin.
4. ADF = approximately cellulose and lignin.
5. Physiological Fuel Value (kcal/gm) = Sum of decimal fractions of protein, fat and carbohydrate (use Nitrogen Free Extract) x 4,9,4 kcal/gm respectively.